

# Classifying semisymmetric cubic graphs of order $20p$

Mohsen SHAHSAVARAN<sup>1</sup>, Mohammad Reza DARAFSHEH<sup>2\*</sup>

(1) School of Mathematics, Statistics, and Computer Science, College of Science, University of Tehran, Tehran, Iran,  
ORCID iD: <https://orcid.org/0000-0002-1416-1388>

(2) School of Mathematics, Statistics, and Computer Science, College of Science, University of Tehran, Tehran, Iran,  
ORCID iD: <https://orcid.org/0000-0002-5539-4989>

---

Received: .201

Accepted/Published Online: .201

Final Version: ..201

---

**Abstract:** A simple graph is called semisymmetric if it is regular and edge transitive but not vertex transitive. In this paper we classify all connected cubic semisymmetric graphs of order  $20p$ ,  $p$  prime.

**Key words:** Edge-transitive graph, vertex-transitive graph, semisymmetric graph, order of a graph, classification of cubic semisymmetric graphs

## 1. Introduction

In this paper all graphs are finite, undirected and simple, i.e. without loops and multiple edges. A graph is called semisymmetric if it is regular and edge-transitive but not vertex-transitive.

The class of semisymmetric graphs was first studied by Folkman [9], who found several infinite families of such graphs and posed eight open problems.

An interesting research problem is to classify connected cubic semisymmetric graphs of various types of orders. In [9], Folkman proved that there are no semisymmetric graphs of order  $2p$  or  $2p^2$  for any prime  $p$ . The classification of semisymmetric graphs of order  $2pq$ , where  $p$  and  $q$  are distinct primes, was given in [7].

For prime  $p$ , cubic semisymmetric graphs of order  $2p^3$  were investigated in [17], in which the authors proved that there is no connected cubic semisymmetric graph of order  $2p^3$  for any prime  $p \neq 3$  and that for  $p = 3$  the only such graph is the Gray graph.

Also connected cubic semisymmetric graphs of orders  $4p^3$ ,  $6p^2$ ,  $6p^3$ ,  $8p^2$ ,  $8p^3$ ,  $10p^3$ ,  $18p^n$  ( $n \geq 1$ ) have been classified in [1, 2, 8, 11, 13, 21].

In this paper we investigate connected cubic semisymmetric graphs of order  $20p$  for all primes  $p$ . Note that for orders like  $4p$ ,  $6p$ ,  $10p$  and  $14p$  which are of the form  $2qp$  for some fixed prime  $q$ , the problem of classifying such graphs follows from the general result of [7].

We prove that if  $\Gamma$  is a connected cubic semisymmetric graph of order  $20p$ ,  $p$  prime, then  $p = 11$  and  $\Gamma$  is isomorphic to a known graph. We go beyond however and prove that there is no connected cubic  $G$ -semisymmetric graph of order  $20p$ , for any prime  $p \neq 2, 11$ . This will put us near the classification of all connected cubic  $G$ -semisymmetric graphs of order  $20p$ : if there is any such graph, then its order must be either

---

\*Correspondence: darafsheh@ut.ac.ir

2010 *AMS Mathematics Subject Classification*: 05E18, 20D60, 05C25, 20B25

1 40 or 220.

2 **2. Preliminaries**

3 In this paper the symmetric and alternating groups of degree  $n$ , the dihedral group of order  $2n$  and the cyclic  
 4 group of order  $n$  are respectively denoted by  $\mathbb{S}_n, \mathbb{A}_n, D_{2n}, \mathbb{Z}_n$ . If  $G$  is a group and  $H \leq G$ , then  $\text{Aut}(G), G',$   
 5  $Z(G), C_G(H)$  and  $N_G(H)$  denote respectively the group of automorphisms of  $G$ , the commutator subgroup  
 6 of  $G$ , the center of  $G$ , the centralizer and the normalizer of  $H$  in  $G$ . We also write  $H \trianglelefteq^c G$  to denote  $H$  is a  
 7 characteristic subgroup of  $G$ . If  $H \trianglelefteq^c K \trianglelefteq G$ , then  $H \trianglelefteq G$ . For a prime  $p$  dividing the order of a finite group  
 8  $G, O_p(G)$  will denote the largest normal  $p$ -subgroup of  $G$ . It is easy to verify that  $O_p(G) \trianglelefteq^c G$ .

9 For a group  $G$  and a nonempty set  $\Omega$ , an action of  $G$  on  $\Omega$  is a function  $(g, \omega) \rightarrow g.\omega$  from  $G \times \Omega$  to  
 10  $\Omega$ , where  $1.\omega = \omega$  and  $g.(h.\omega) = (gh).\omega$ , for every  $g, h \in G$  and every  $\omega \in \Omega$ . We write  $g\omega$  instead of  $g.\omega$ , if  
 11 there is no fear of ambiguity. For  $\omega \in \Omega$ , the stabilizer of  $\omega$  in  $G$  is defined as  $G_\omega = \{g \in G : g\omega = \omega\}$ . The  
 12 action is called *semiregular* if the stabilizer of each element in  $\Omega$  is trivial; it is called *regular* if it is semiregular  
 13 and transitive.

14 For any two groups  $G$  and  $H$  and any homomorphism  $\varphi : H \rightarrow \text{Aut}(G)$  the *external semidirect product*  
 15  $G \rtimes_\varphi H$  is defined as the group whose underlying set is the cartesian product  $G \times H$  and whose binary operation  
 16 is defined as  $(g_1, h_1)(g_2, h_2) = (g_1\varphi(h_1)(g_2), h_1h_2)$ . If  $\varphi(h) = 1$  for each  $h \in H$ , then the semidirect product  
 17 will coincide with the usual direct product. If  $G = NK$  where  $N \trianglelefteq G, K \leq G$  and  $N \cap K = 1$ , then  $G$  is said  
 18 to be the *internal semidirect product* of  $N$  and  $K$ . These two concepts are in fact equivalent in the sense that  
 19 there is some homomorphism  $\varphi : K \rightarrow \text{Aut}(N)$  where  $G \simeq N \rtimes_\varphi K$ .

20 The dihedral group  $D_{2n}$  is defined as

21 
$$D_{2n} = \langle a, b | a^n = b^2 = 1, b^{-1}ab = a^{-1} \rangle$$

22 So  $D_{2n} = \{a^i | i = 0, \dots, n - 1\} \cup \{ba^i | i = 0, \dots, n - 1\}$ . All the elements of the form  $ba^i$  are of order 2.

23 Let  $\Gamma$  be a graph. For two vertices  $u$  and  $v$ , we write  $u \sim v$  to denote  $u$  is adjacent to  $v$ . If  $u \sim v$ ,  
 24 then each of the ordered pairs  $(u, v)$  and  $(v, u)$  is called an *arc*. The set of all vertices adjacent to a vertex  
 25  $u$  is denoted by  $\Gamma(u)$ . The degree or valency of  $u$  is  $|\Gamma(u)|$ . We call  $\Gamma$  *regular* if all of its vertices have the  
 26 same valency. The vertex set, the edge set, the arc set and the set of all automorphisms of  $\Gamma$  are respectively  
 27 denoted by  $V(\Gamma), E(\Gamma), \text{Arc}(\Gamma)$  and  $\text{Aut}(\Gamma)$ . If  $\Gamma$  is a graph and  $N \trianglelefteq \text{Aut}(\Gamma)$ , then  $\Gamma_N$  will denote a simple  
 28 undirected graph whose vertices are the orbits of  $N$  in its action on  $V(\Gamma)$ , and where two vertices  $Nu$  and  $Nv$   
 29 are adjacent if and only if  $u \sim nv$  in  $\Gamma$ , for some  $n \in N$ .

30 Let  $\Gamma_c$  and  $\Gamma$  be two graphs. Then  $\Gamma_c$  is said to be a *covering graph* for  $\Gamma$  if there is a surjection  
 31  $f : V(\Gamma_c) \rightarrow V(\Gamma)$  which preserves adjacency and for each  $u \in V(\Gamma_c)$ , the restricted function  $f|_{\Gamma_c(u)} :$   
 32  $\Gamma_c(u) \rightarrow \Gamma(f(u))$  is a one to one correspondence.  $f$  is called a *covering projection*. Clearly, if  $\Gamma$  is bipartite,  
 33 then so is  $\Gamma_c$ . For each  $u \in V(\Gamma)$ , the *fibre* on  $u$  is defined as  $\text{fib}_u = f^{-1}(u)$ . The following important set is  
 34 a subgroup of  $\text{Aut}(\Gamma_c)$  and is called the *group of covering transformations* for  $f$ :

35 
$$CT(f) = \{\sigma \in \text{Aut}(\Gamma_c) | \forall u \in V(\Gamma), \sigma(\text{fib}_u) = \text{fib}_u\}$$

36 It is known that  $K = CT(f)$  acts semiregularly on each fibre [14]. If this action is regular, then  $\Gamma_c$  is said to  
 37 be a *regular K-cover* of  $\Gamma$ .

38 Let  $X \leq \text{Aut}(\Gamma)$ . Then  $\Gamma$  is said to be *X-vertex transitive, X-edge transitive* or *X-arc transitive* if  
 39  $X$  acts transitively on  $V(\Gamma), E(\Gamma)$  or  $\text{Arc}(\Gamma)$  respectively. The graph  $\Gamma$  is called *X-semisymmetric* if it is

1 regular and  $X$ -edge transitive but not  $X$ -vertex transitive. Also  $\Gamma$  is called  $X$ -symmetric if it is  $X$ -vertex  
 2 transitive and  $X$ -arc transitive. For  $X = \text{Aut}(\Gamma)$ , we omit  $X$  and simply talk about  $\Gamma$  being edge transitive,  
 3 vertex transitive, symmetric or semisymmetric. As an example,  $\Gamma = K_{3,3}$ , the complete bipartite graph on 6  
 4 vertices, is not semisymmetric but it is  $X$ -semisymmetric for some  $X \leq \text{Aut}(\Gamma)$ .

5 An  $X$ -edge transitive but not  $X$ -vertex transitive graph is necessarily bipartite, where the two partites  
 6 are the orbits of the action of  $X$  on  $V(\Gamma)$ . If  $\Gamma$  is regular, then the two partite sets have equal cardinality. So  
 7 an  $X$ -semisymmetric graph is bipartite such that  $X$  is transitive on each partite but  $X$  carries no vertex from  
 8 one partite set to the other.

9 According to [5], if there is a unique known cubic semisymmetric graph of order  $n$ , then it is denoted by  
 10 **Sn**. The symmetric counterpart of **Sn** is denoted by **Fn** ([6]). There are only two symmetric cubic graphs of  
 11 order 20 which are denoted by **F20A** and **F20B**. Only **F20B** is bipartite ([6]).

12 Any minimal normal subgroup of a finite group, is the internal direct product of isomorphic copies of a  
 13 simple group.

14 A finite group  $G$  is called a  $K_n$ -group if its order has exactly  $n$  distinct prime divisors, where  $n \in \mathbb{N}$ .  
 15 The following two results determine all simple  $K_3$ -groups and  $K_4$ -groups [3, 12, 19, 24].

16 **Theorem 2.1** (i) If  $G$  is a simple  $K_3$ -group, then  $G$  is isomorphic to one of the following groups:  $\mathbb{A}_5, \mathbb{A}_6,$   
 17  $L_2(7), L_2(2^3), L_2(17), L_3(3), U_3(3), U_4(2)$ .

18 (ii) If  $G$  is a simple  $K_4$ -group, then  $G$  is isomorphic to one of the following groups:

19 (1)  $\mathbb{A}_7, \mathbb{A}_8, \mathbb{A}_9, \mathbb{A}_{10}, M_{11}, M_{12}, J_2, L_2(2^4), L_2(5^2), L_2(7^2), L_2(3^4), L_2(97), L_2(3^5), L_2(577), L_3(2^2),$   
 20  $L_3(5), L_3(7), L_3(2^3), L_3(17), L_4(3), U_3(2^2), U_3(5), U_3(7), U_3(2^3), U_3(3^2), U_4(3), U_5(2), S_4(2^2),$   
 21  $S_4(5), S_4(7), S_4(3^2), S_6(2), O_8^+(2), G_2(3), Sz(2^3), Sz(2^5), {}^3D_4(2), {}^2F_4(2)'$ ;

22 (2)  $L_2(r)$  where  $r$  is a prime,  $r^2 - 1 = 2^a \cdot 3^b \cdot s$ ,  $s > 3$  is a prime,  $a, b \in \mathbb{N}$ ;

23 (3)  $L_2(2^m)$  where  $m, 2^m - 1, \frac{2^m+1}{3}$  are primes greater than 3;

24 (4)  $L_2(3^m)$  where  $m, \frac{3^m+1}{4}$  and  $\frac{3^m-1}{2}$  are odd primes.

25 **Proposition 2.2** ([18], Theorem 9.1.2) Let  $G$  be a finite group and  $N \trianglelefteq G$ . If  $|N|$  and  $|\frac{G}{N}|$  are relatively  
 26 prime, then  $G$  has a subgroup  $H$  such that  $G = NH$  and  $N \cap H = 1$  (therefore  $G$  is the internal semidirect  
 27 product of  $N$  and  $H$ ).

28 An immediate consequence of the following theorem of Burnside is that the order of every nonabelian  
 29 simple group is divisible by at least 3 distinct primes.

30 **Theorem 2.3** [20] For any two distinct primes  $p$  and  $q$  and any two nonnegative integers  $a$  and  $b$ , every  
 31 finite group of order  $p^a q^b$  is solvable.

32 In the following theorem, the inverse of a pair  $(a, b)$ , is meant to be  $(b, a)$ . Also for each  $i, A_i, B_i, C_i$   
 33 and  $D_i$  are certain groups of order  $i$  with known structures. We will not need their structures.

1 **Theorem 2.4** [10] *If  $\Gamma$  is a connected cubic  $X$ -semisymmetric graph, then the order of the stabilizer of any*  
 2 *vertex is of the form  $2^r \cdot 3$  for some  $0 \leq r \leq 7$ . More precisely, if  $\{u, v\}$  is any edge of  $\Gamma$ , then the pair*  
 3  *$(X_u, X_v)$  can only be one of the following fifteen pairs or their inverses:*

4  $(\mathbb{Z}_3, \mathbb{Z}_3), (\mathbb{S}_3, \mathbb{S}_3), (\mathbb{S}_3, \mathbb{Z}_6), (D_{12}, D_{12}), (D_{12}, \mathbb{A}_4), (\mathbb{S}_4, D_{24}), (\mathbb{S}_4, \mathbb{Z}_3 \rtimes D_8), (\mathbb{A}_4 \times \mathbb{Z}_2, D_{12} \times \mathbb{Z}_2), (\mathbb{S}_4 \times \mathbb{Z}_2, D_8 \times$   
 5  $\mathbb{S}_3), (\mathbb{S}_4, \mathbb{S}_4), (\mathbb{S}_4 \times \mathbb{Z}_2, \mathbb{S}_4 \times \mathbb{Z}_2), (A_{96}, B_{96}), (A_{192}, B_{192}), (C_{192}, D_{192}), (A_{384}, B_{384}).$

6 **Proposition 2.5** [17] *Let  $\Gamma$  be a connected cubic  $X$ -semisymmetric graph for some  $X \leq \text{Aut}(\Gamma)$ ; then either*  
 7  *$\Gamma \simeq K_{3,3}$ , the complete bipartite graph on 6 vertices, or  $X$  acts faithfully on each of the bipartition sets of  $\Gamma$ .*

8 **Theorem 2.6** [15] *Let  $\Gamma$  be a connected cubic  $X$ -semisymmetric graph. Let  $\{U, W\}$  be a bipartition for  $\Gamma$*   
 9 *and assume  $N \trianglelefteq X$ . If the actions of  $N$  on both  $U$  and  $W$  are intransitive, then  $N$  acts semiregularly on both*  
 10  *$U$  and  $W$ ,  $\Gamma_N$  is  $\frac{X}{N}$ -semisymmetric, and  $\Gamma$  is a regular  $N$ -covering of  $\Gamma_N$ .*

11 This theorem has a nice result. For every normal subgroup  $N \trianglelefteq X$  either  $N$  is transitive on at least one partite  
 12 set or it is intransitive on both partite sets. In the former case, the order of  $N$  is divisible by  $|U| = |W|$ . In  
 13 the latter case, according to Theorem 2.6, the induced action of  $N$  on both  $U$  and  $W$  is semiregular and hence  
 14 the order of  $N$  divides  $|U| = |W|$ . So we have the following handy corollary.

15 **Corollary 2.7** *If  $\Gamma$  is a connected cubic  $X$ -semisymmetric graph with  $\{U, W\}$  as a bipartition and  $N \trianglelefteq X$ ,*  
 16 *then either  $|N|$  divides  $|U|$  or  $|U|$  divides  $|N|$ .*

17 Following [10](see also [16]) the coset graph  $C(G; H_0, H_1)$  of a group  $G$  with respect to finite subgroups  
 18  $H_0$  and  $H_1$  is a bipartite graph with  $\{H_0g | g \in G\}$  and  $\{H_1g | g \in G\}$  as its bipartition sets of vertices where  
 19  $H_0g$  is adjacent to  $H_1g'$  whenever  $H_0g \cap H_1g' \neq \emptyset$ . The following proposition may be extracted from [10]:

20 **Proposition 2.8** *Let  $G$  be a finite group and  $H_0, H_1 \leq G$ . The coset graph  $C(G; H_0, H_1)$  has the following*  
 21 *properties:*

22 (i)  $C(G; H_0, H_1)$  is regular of valency  $d$  if and only if  $H_0 \cap H_1$  has index  $d$  in both  $H_0$  and  $H_1$ .

23 (ii)  $C(G; H_0, H_1)$  is connected if and only if  $G = \langle H_0, H_1 \rangle$ .

24 (iii)  $G$  acts on  $C(G; H_0, H_1)$  by right multiplication. Moreover this action is faithful if and only if  $\text{Core}_G(H_0 \cap$   
 25  $H_1) = 1$ .

26 (iv) In the case when the action of  $G$  is faithful, the coset graph  $C(G; H_0, H_1)$  is  $G$ -semisymmetric.

27 **Proposition 2.9** [16] *Let  $\Gamma$  be a regular graph and  $G \leq \text{Aut}(\Gamma)$ . If  $\Gamma$  is  $G$ -semisymmetric, then  $\Gamma$  is*  
 28 *isomorphic to the coset graph  $C(G; G_u, G_v)$  where  $u$  and  $v$  are adjacent vertices.*

### 29 3. Main Results

30 Our goal in this paper is to fully classify connected cubic semisymmetric graphs of order  $20p$ . We also derive a  
 31 very restrictive necessary condition for the existence of connected cubic  $G$ -semisymmetric graphs of order  $20p$ .  
 32 We prove the following important result. Part (i) is a full classification whereas part (ii) is only a necessary  
 33 condition.

**Theorem 3.1** *Let  $p$  be a prime.*

(i) *If  $\Gamma$  is a connected cubic semisymmetric graph of order  $20p$ , then  $p = 11$  and  $\Gamma \simeq S220$ .*

(ii) *If  $\Gamma$  is a connected cubic  $G$ -semisymmetric graph of order  $20p$  for some  $G \leq \text{Aut}(\Gamma)$ , then  $p = 2$  or  $11$ .*

To prove the main theorem, we need some lemmas.

**Lemma 3.2** *The only simple  $K_4$ -groups whose orders are of the form  $2^i \cdot 3 \cdot 5 \cdot p$  for some prime  $p > 5$  and some  $1 \leq i \leq 8$ , are the following three projective special linear groups:  $L_2(2^4)$ ,  $L_2(11)$  and  $L_2(31)$ .*

**Proof** Considering the powers of primes, there is no possibility for such a group in sub-item (4) of item (ii) of Theorem 2.1. By inspecting orders of groups in sub-item (1), the only group of the desired form is  $L_2(2^4)$ . As for sub-item (3), let  $L_2(2^m)$  be a group of order  $2^i \cdot 3 \cdot 5 \cdot p$ ; then

$$2^m \cdot 3 \cdot (2^m - 1) \cdot \left(\frac{2^m+1}{3}\right) = 2^i \cdot 3 \cdot 5 \cdot p$$

where  $m$ ,  $2^m - 1$  and  $\frac{2^m+1}{3}$  are all primes according to Theorem 2.1. This equation has no answer as neither  $2^m - 1$  nor  $\frac{2^m+1}{3}$  could be equal to 5. Finally consider groups  $L_2(r)$  in sub-item (2). If for odd prime  $r$  and for prime  $s > 3$ , we have  $r^2 - 1 = 2^a \cdot 3^b \cdot s$  and

$$2^{a-1} \cdot 3^b \cdot s \cdot r = 2^i \cdot 3 \cdot 5 \cdot p,$$

then  $b = 1$ ,  $a - 1 = i$  and either  $s = 5$  or  $r = 5$ . The equality  $r = 5$  is not possible, since  $L_2(5)$  is not a  $K_4$ -group. Also if  $s = 5$ , then the equation  $r^2 - 1 = 2^a \cdot 3 \cdot 5$  gives us only two solutions  $r = 11, 31$  when  $a$  spans integers  $2, 3, \dots, 9$ . □

**Lemma 3.3** *Let  $p > 11$  be a prime and  $p \neq 17, 31$ . If  $\Gamma$  is a connected cubic  $G$ -semisymmetric graph of order  $20p$ , then  $G$  has a normal Sylow  $p$ -subgroup.*

**Proof** Take  $\{U, W\}$  to be a bipartition for  $\Gamma$ . Then  $|U| = |W| = 10p$ . For  $u \in U$  according to Theorem 2.4,  $|G_u| = 2^r \cdot 3$  for some  $0 \leq r \leq 7$ . Due to transitivity of  $G$  on  $U$ , the equality  $[G : G_u] = |U|$  holds which yields  $|G| = 2^{r+1} \cdot 3 \cdot 5 \cdot p$ . If  $G$  does not have a normal Sylow  $p$ -subgroup, then  $O_p(G) = 1$ . We derive a contradiction out of this.

Suppose  $G$  has a normal subgroup  $M$  of order 10. Due to its order,  $M$  is intransitive on the partite sets and according to Theorem 2.6, the quotient graph  $\Gamma_M$  is  $\frac{G}{M}$ -semisymmetric with a bipartition  $\{U_M, W_M\}$  where  $|U_M| = |W_M| = p$  and  $|\frac{G}{M}| = 2^r \cdot 3 \cdot p$ .

Let  $\frac{K}{M}$  be a minimal normal subgroup of  $\frac{G}{M}$ . If  $\frac{K}{M}$  is unsolvable, it must be simple of order  $2^i \cdot 3 \cdot p$  for some  $i$ . So  $\frac{K}{M} \simeq A_5$  or  $L_2(7)$ . However these are not possible since  $p > 11$ . Now if  $\frac{K}{M}$  is solvable and hence elementary abelian, then by Corollary 2.7, its order must be  $p$  implying  $|K| = 10p$ . The Sylow  $p$ -subgroup of  $K$  is normal and hence characteristic in  $K$ . Therefore it is normal in  $G$ , contradicting the assumption that  $O_p(G) = 1$ . So  $O_p(G) = 1$  implies that  $G$  does not have a normal subgroup of order 10.

Next, let  $N \simeq T^k$  be a minimal normal subgroup of  $G$ , where  $T$  is simple. If  $T$  is nonabelian, then  $k = 1$  and  $N = T$  since the powers of 3 and 5 in  $|G|$  equal 1. According to Corollary 2.7, either  $|N|$  divides  $|U| = 10p$  or  $10p$  divides  $|N|$ .

If  $|N|$  divides  $10p$ , then  $|N| = 2 \cdot 5 \cdot p$ , since  $|N|$  should be divisible by at least three distinct primes (Theorem 2.3). But there is no simple  $K_3$ -group of order  $2 \cdot 5 \cdot p$  according to part (i) of Theorem 2.1. So  $10p$  divides  $|N|$ . Since the order of every simple  $K_3$ -group is divisible by 3,  $N$  must be a simple  $K_4$ -group whose order is of the form  $2^i \cdot 3 \cdot 5 \cdot p$ . According to Lemma 3.2,  $N \simeq L_2(2^4)$ ,  $L_2(11)$  or  $L_2(31)$  corresponding to  $p = 17, 11$  and  $31$  respectively. But these cases are ruled out in the statement of the Lemma.

Now suppose  $T$  is abelian and hence  $N$  would be elementary abelian. It follows from Corollary 2.7, that  $|N|$  divides  $10p$  and so  $|N| = 2, 5$  or  $p$ . Certainly  $|N| = p$  contradicts the assumption on  $O_p(G)$ . In the remaining two cases  $\Gamma_N$  would itself be a connected cubic  $\frac{G}{N}$ -semisymmetric graph of order  $\frac{20p}{|N|}$ . Take  $\{U_N, W_N\}$  to be the bipartition for  $\Gamma_N$ . Also let  $\frac{M}{N}$  be a minimal normal subgroup of  $\frac{G}{N}$ .

If  $N \simeq \mathbb{Z}_2$ , then  $|\frac{G}{N}| = 2^r \cdot 3 \cdot 5 \cdot p$  and  $|U_N| = |W_N| = 5p$ . If  $\frac{M}{N}$  is unsolvable, then it must be a simple  $K_4$ -group whose order is of the form  $2^i \cdot 3 \cdot 5 \cdot p$ . It follows from Lemma 3.2, that  $p = 17, 11$  or  $31$  which are ruled out by our assumption on  $p$ . On the other hand if  $\frac{M}{N}$  is solvable, then its order should divide  $|U_N| = 5p$  and hence  $|\frac{M}{N}| = 5$  or  $p$ . If  $|\frac{M}{N}| = 5$ , then  $|M| = 10$  which is not possible (as we showed at the beginning of the proof), and if  $|\frac{M}{N}| = p$ , then  $|M| = 2p$  which contradicts our assumption on  $O_p(G)$  since a Sylow  $p$ -subgroup of  $M$  would be characteristic in  $M$  and so would be normal in  $G$ .

Now if  $N \simeq \mathbb{Z}_5$ , then  $|\frac{G}{N}| = 2^{r+1} \cdot 3 \cdot p$  and  $|U_N| = |W_N| = 2p$ . In this case if  $\frac{M}{N}$  is unsolvable, it would be a simple group of order  $2^i \cdot 3 \cdot p$  for some  $i$  and hence according to Theorem 2.1,  $\frac{M}{N} \simeq \mathbb{A}_5$  or  $L_2(7)$  implying  $p = 5$  or  $7$ . This is in contradiction to our assumption on  $p$ . On the other hand if  $\frac{M}{N}$  is solvable, then like before, we conclude that  $|\frac{M}{N}| = 2$  or  $p$  which again lead to contradictions as in the previous case.  $\square$

**Lemma 3.4** *Let  $p > 11$  be a prime and  $p \neq 17, 31$ . Suppose  $\Gamma$  is a connected cubic  $G$ -semisymmetric graph of order  $20p$ . Let  $M$  be the Sylow  $p$ -subgroup of  $G$ . If  $\frac{G}{M} \simeq H$ , then*

- (1) *For each vertex  $u$  the stabilizer  $G_u$  is isomorphic to a subgroup of  $H$ .*
- (2)  *$G \simeq M \rtimes_{\varphi} H$  for some homomorphism  $\varphi : H \rightarrow \text{Aut}(M)$ .*

**Proof** For each vertex  $u$  of  $\Gamma$ ,  $MG_u \leq G$ . Therefore  $G_u \simeq \frac{G_u}{M \cap G_u} \simeq \frac{MG_u}{M} \leq \frac{G}{M}$ . This proves (1). Now since obviously the orders of  $M$  and  $\frac{G}{M}$  are coprime, it follows from Proposition 2.2, that  $G = MK$  for some subgroup  $K \leq G$  where  $M \cap K = 1$ . So  $G$  is the internal semidirect product of  $M$  and  $K$  and hence it is isomorphic to the external semidirect product of  $M$  and  $K$ ; i.e.  $G \simeq M \rtimes_{\psi} K$  for some  $\psi : K \rightarrow \text{Aut}(M)$ . Since  $H \simeq \frac{G}{M} = \frac{MK}{M} \simeq \frac{K}{M \cap K} \simeq K$ , we can write  $G \simeq M \rtimes_{\varphi} H$  for some  $\varphi : H \rightarrow \text{Aut}(M)$ .  $\square$

**Lemma 3.5** *Let  $p > 11$  be a prime and  $p \neq 17, 31$ . If  $\Gamma$  is a connected cubic  $G$ -semisymmetric graph of order  $20p$  and if  $M$  is the Sylow  $p$ -subgroup of  $G$ , then  $\frac{G}{M}$  cannot be isomorphic to  $\mathbb{A}_5$ .*

**Proof** Suppose on the contrary, that  $\frac{G}{M} \simeq \mathbb{A}_5$ . Then for any vertex  $u$  from  $[G : G_u] = 10p$  we obtain  $|G_u| = 6$  and hence  $G_u \simeq \mathbb{Z}_6$  or  $\mathbb{S}_3$ . By Lemma 3.4,  $G_u \leq \mathbb{A}_5$ . Since  $\mathbb{A}_5$  does not have elements of order 6, we conclude that  $G_u \simeq \mathbb{S}_3$ . Also according to Lemma 3.4,  $G \simeq M \rtimes_{\varphi} \mathbb{A}_5$ . There are only two possibilities for the kernel of  $\varphi : \mathbb{A}_5 \rightarrow \text{Aut}(M)$ .

(a) If  $\ker(\varphi) = 1$ , then  $\mathbb{A}_5$  is isomorphic to a subgroup of  $\text{Aut}(M) \simeq \text{Aut}(\mathbb{Z}_p) \simeq \mathbb{Z}_{p-1}$  which is obviously not the case.

(b) If  $\ker(\varphi) = \mathbb{A}_5$ , then  $\varphi$  is the trivial homomorphism and so  $G \simeq M \times \mathbb{A}_5$ . Since  $\Gamma$  is  $G$ -semisymmetric, according to Proposition 2.9,  $\Gamma$  is isomorphic to  $C(G; G_u, G_v)$  where  $u$  and  $v$  are two adjacent vertices in  $\Gamma$ . As  $\Gamma$  is connected, according to Proposition 2.8, we must have  $G = \langle G_u, G_v \rangle$ . In view of  $G_u \simeq G_v \simeq \mathbb{S}_3$ , this means that  $M \times \mathbb{A}_5$  is generated by two of its subgroups, say  $H$  and  $K$ , both isomorphic to  $\mathbb{S}_3$ . Now for each element  $(m, a) \in H$  we have  $(m, a)^6 = 1$  which means  $m^6 = 1$  in  $M$ . As  $|M| = p > 31$ , we conclude  $m = 1$ . Therefore the first component of each element of  $H$  (and similarly for  $K$ ) equals 1. Consequently the first component of each element in  $M \times \mathbb{A}_5 = \langle H, K \rangle$  equals 1 which is a contradiction.  $\square$

Consider a semidirect product  $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  where  $\varphi : \mathbb{S}_5 \rightarrow \text{Aut}(\mathbb{Z}_p)$  is a homomorphism. Let  $H \leq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  and  $H \simeq D_{12}$  or  $\mathbb{A}_4$ . We call  $H$  of **type A** if all the elements of  $H$  have their second component in  $\mathbb{A}_5$ . We also call  $H$  of **type D** if there is at least one element in  $H$  whose second component is not in  $\mathbb{A}_5$ . Also for any  $x \in \mathbb{Z}_p$  and any  $g, h \in \mathbb{S}_5$  we define two subsets  $R_{x,g,h}, S_{x,g,h} \subset \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  as follows:

$$R_{x,g,h} = \{(1, 1), (x, g), (1, g^2), (x, g^3), (1, g^4), (x, g^5), (1, h), (x, hg), (1, hg^2), (x, hg^3), (1, hg^4), (x, hg^5)\}$$

and

$$S_{x,g,h} = \{(1, 1), (x, g), (1, g^2), (x, g^3), (1, g^4), (x, g^5), (x, h), (1, hg), (x, hg^2), (1, hg^3), (x, hg^4), (1, hg^5)\}.$$

As we will see later, these two subsets are sometimes subgroups of  $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$ .

The group  $D_{12} = \langle a, b \mid a^6 = b^2 = 1, b^{-1}ab = a^{-1} \rangle$  has exactly three Sylow 2-subgroups, all isomorphic to  $\mathbb{Z}_2 \times \mathbb{Z}_2$ , which are listed below:

$$P_1 = \{1, a^3, b, ba^3\}, P_2 = \{1, a^3, ba, ba^4\}, P_3 = \{1, a^3, ba^2, ba^5\}.$$

**Lemma 3.6** *Let  $p > 3$  be a prime and let  $\varphi : \mathbb{S}_5 \rightarrow \text{Aut}(\mathbb{Z}_p)$  be a homomorphism where  $\ker(\varphi) = \mathbb{A}_5$ . Let  $H \leq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$ .*

(i) *If  $H \simeq \mathbb{A}_4$  then  $H$  is of type A and if  $H \simeq D_{12}$  then  $H$  is of type D.*

(ii) *Moreover if  $H \simeq D_{12}$ , then there are some  $x \in \mathbb{Z}_p$ , some  $g, g' \notin \mathbb{A}_5$  and some  $h \in \mathbb{A}_5$  where  $H = R_{x,g,h}$  or  $H = S_{x,g,g'}$ .*

**Proof** The image of  $\varphi$  is isomorphic to  $\frac{\mathbb{S}_5}{\mathbb{A}_5} \simeq \mathbb{Z}_2$ . So there is some  $F \in \text{Aut}(\mathbb{Z}_p)$  of order 2 for which  $\varphi(x) = 1$  for all  $x \in \mathbb{A}_5$  and  $\varphi(x) = F$  for any  $x \notin \mathbb{A}_5$ . For any two elements  $(x, g)$  and  $(y, h)$  from  $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  the multiplication  $(x, g)(y, h)$  equals  $(xy, gh)$  if  $g \in \mathbb{A}_5$  and equals  $(xF(y), gh)$  if  $g \notin \mathbb{A}_5$ . It is easy to see that for any positive integer  $n$  if  $g \in \mathbb{A}_5$ , then  $(x, g)^n = (x^n, g^n)$  for all  $x \in \mathbb{Z}_p$ , and if  $g \notin \mathbb{A}_5$ , then  $(x, g)^{2n} = (x^n F(x^n), g^{2n})$  and  $(x, g)^{2n+1} = (x^{n+1} F(x^n), g^{2n+1})$  for all  $x \in \mathbb{Z}_p$ .

Now to prove part (i), it suffices to prove that for a subgroup  $H \leq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  which is isomorphic to either  $D_{12}$  or  $\mathbb{A}_4$ , both of the following statements are true:

- if  $H$  is of type A, then  $H \simeq \mathbb{A}_4$
- if  $H$  is of type D, then  $H \simeq D_{12}$

1 Let  $H \leq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  and  $H \simeq D_{12}$  or  $\mathbb{A}_4$ . If  $K := \{(x, g) \in H | g \in \mathbb{A}_5\}$  then  $K \leq H$ . For the  
 2 homomorphism  $f : K \rightarrow \mathbb{Z}_p$  defined by  $f(x, g) = x$ , the isomorphism  $\frac{K}{\ker(f)} \simeq \text{Im}(f)$  implies that  $|\frac{K}{\ker(f)}|$   
 3 divides both 12 and  $p$  and hence  $K = \ker(f)$ . Therefore for each  $(x, g) \in H$  if  $g \in \mathbb{A}_5$ , then  $x = 1$ .

4 It follows immediately that if  $H$  is of type A, then the first component of each element of  $H$  equals 1  
 5 and hence  $H$  is isomorphic to a subgroup of  $\mathbb{A}_5$ . As  $\mathbb{A}_5$  has no element of order 6,  $H$  cannot be isomorphic  
 6 to  $D_{12}$  and so  $H \simeq \mathbb{A}_4$ .

7 Now suppose  $H$  is of type D. For two arbitrary elements  $(x, g), (y, h) \in H$  with  $g, h \notin \mathbb{A}_5$ , we have  
 8  $(xF(x), g^2) = (x, g)^2 \in H$  and  $(yF(x), hg) = (y, h)(x, g) \in H$ . Since  $g^2$  and  $hg$  are in  $\mathbb{A}_5$ , the first components  
 9 must equal 1; i.e.  $xF(x) = 1$  and  $yF(x) = 1$  which imply  $x = y$ . In other words, for any pair of elements  
 10  $(x, g) \in H$  and  $(y, h) \in H$  with  $g, h \notin \mathbb{A}_5$  we must have  $x = y$ . There are always elements in  $H$  whose second  
 11 component lies in  $\mathbb{A}_5$  and hence their first component is 1. Therefore we can write

$$H = \{(x, g_1), (x, g_2), \dots, (x, g_n), (1, h_1), \dots, (1, h_m)\} \tag{3.1}$$

12 where  $n + m = 12$  and where  $g_1, \dots, g_n \notin \mathbb{A}_5$  and  $h_1, \dots, h_m \in \mathbb{A}_5$ . It also follows that for this specific  $x$ ,  
 13  $F(x) = x^{-1}$ . Let

$$\overline{H} = \{(1, h_1), \dots, (1, h_m)\}, H_1 = \{h_1, \dots, h_m\};$$

15 then  $H_1 \simeq \overline{H}$ ,  $\overline{H} \leq H$  and  $H_1 \leq \mathbb{A}_5$ . Multiplying all the elements of  $H$  from equation 3.1, by  $(x, g_t)$  for an  
 16 arbitrary  $t$ , we again obtain  $H$ . Therefore

$$H = \{(1, g_t g_1), (1, g_t g_2), \dots, (1, g_t g_n), (x, g_t h_1), \dots, (x, g_t h_m)\}. \tag{3.2}$$

17 Comparing the equalities 3.1 and 3.2 and by taking into account that  $g_t g_i \in \mathbb{A}_5$  for  $i = 1, \dots, n$  and  $g_t h_j \notin \mathbb{A}_5$   
 18 for  $j = 1, \dots, m$ , it follows that

$$\{g_t h_1, \dots, g_t h_m\} = \{g_1, \dots, g_n\}.$$

20 Therefore  $m = n = 6$  and so  $|\overline{H}| = 6$ . Since  $\mathbb{A}_4$  does not have a subgroup of order 6, we conclude that  
 21  $H \simeq D_{12}$ .

22 We now proceed to prove part (ii). So let  $D_{12} \simeq H \leq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$ . According to part (i)  $H$  is of type D.  
 23 We continue to use the notations invented in the proof of part (i). The group  $D_{12}$  has only two subgroups of  
 24 order 6, namely  $\mathbb{Z}_6$  and  $\mathbb{S}_3$ . Since  $\overline{H} \leq H$  and  $|\overline{H}| = 6$ , we have  $\overline{H} \simeq \mathbb{Z}_6$  or  $\mathbb{S}_3$ . Since  $\overline{H} \simeq H_1 \leq \mathbb{A}_5$  and  $\mathbb{A}_5$   
 25 does not have elements of order 6, it follows that  $\overline{H}$  cannot be isomorphic to  $\mathbb{Z}_6$  and hence  $\overline{H} \simeq \mathbb{S}_3$ . Also as  
 26  $H \simeq D_{12}$ , we can write  $H = \{a^i | i = 0, \dots, 5\} \cup \{ba^i | i = 0, \dots, 5\}$ . As  $\overline{H} \simeq \mathbb{S}_3$  does not have any element of  
 27 order 6, we must have  $a \in H - \overline{H}$ ; i.e.  $a = (x, g)$  for some  $g \notin \mathbb{A}_5$  (see equation 3.1). As for  $b$ , there are two  
 28 possible cases; either  $b = (1, h) \in \overline{H}$  or  $b = (x, g') \in H - \overline{H}$ .

29 If  $b = (1, h)$ ,  $h \in \mathbb{A}_5$ , then

$$30 H = \{(x, g)^i | i = 0, \dots, 5\} \cup \{(1, h)(x, g)^i | i = 0, \dots, 5\} = \{(1, 1), (x, g), (1, g^2), (x, g^3), (1, g^4), (x, g^5),$$

$$31 (1, h), (x, hg), (1, hg^2), (x, hg^3), (1, hg^4), (x, hg^5)\} = R_{x, g, h}.$$

32 Also if  $b = (x, g')$ ,  $g' \notin \mathbb{A}_5$ , then

$$H = \{(x, g)^i | i = 0, \dots, 5\} \cup \{(x, g')(x, g)^i | i = 0, \dots, 5\} = \{(1, 1), (x, g), (1, g^2), (x, g^3), (1, g^4), (x, g^5), (x, g'), (1, g'g), (x, g'g^2), (1, g'g^3), (x, g'g^4), (1, g'g^5)\} = S_{x, g, g'}.$$

□

**Lemma 3.7** *Let  $p > 3$  be a prime. A semidirect product  $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  does not have two subgroups  $U$  and  $V$  with all the following properties:*

- 1)  $(U, V) \simeq (D_{12}, D_{12})$  or  $(D_{12}, \mathbb{A}_4)$ ; and
- 2)  $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5 = \langle U, V \rangle$ ; and
- 3)  $U \cap V$  is a common Sylow 2-subgroup of both  $U$  and  $V$ .

**Proof** Let  $\varphi : \mathbb{S}_5 \rightarrow \text{Aut}(\mathbb{Z}_p)$  be a homomorphism. The kernel of  $\varphi$  could not be identity since otherwise  $\mathbb{S}_5$  would be isomorphic to a subgroup of  $\text{Aut}(\mathbb{Z}_p) \simeq \mathbb{Z}_{p-1}$  which is impossible. On the other hand if  $\ker(\varphi) = \mathbb{S}_5$ , then  $\varphi$  is the trivial homomorphism and so  $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5 = \mathbb{Z}_p \times \mathbb{S}_5$ . For any two subgroups  $U, V \leq \mathbb{Z}_p \times \mathbb{S}_5$  both of order 12, the equality  $(x, a)^{12} = 1$  holds for each  $(x, a) \in U \cup V$ . This implies  $x^{12} = 1$  and hence  $x = 1$ , since  $x \in \mathbb{Z}_p$ . Consequently the equality  $\mathbb{Z}_p \times \mathbb{S}_5 = \langle U, V \rangle$  cannot hold.

The only remaining possibility is to have  $\ker(\varphi) = \mathbb{A}_5$ . We assume there are subgroups  $U, V$  with the desired properties and reach a contradiction. So  $U \simeq D_{12}$  and hence according to Lemma 3.6, there are some  $x \in \mathbb{Z}_p$ , some  $g, k \notin \mathbb{A}_5$  and some  $h \in \mathbb{A}_5$  where  $U = R_{x, g, h}$  or  $U = S_{x, g, k}$ . If  $U = R_{x, g, h}$ , then all the Sylow 2-subgroups of  $U$  are as follows:

$$\begin{aligned} RP_{x, g, h}^1 &= \{(1, 1), (x, g^3), (1, h), (x, hg^3)\}, \\ RP_{x, g, h}^2 &= \{(1, 1), (x, g^3), (x, hg), (1, hg^4)\}, \\ RP_{x, g, h}^3 &= \{(1, 1), (x, g^3), (1, hg^2), (x, hg^5)\} \end{aligned}$$

and if  $U = S_{x, g, k}$ , then all the Sylow 2-subgroups of  $U$  are as follows:

$$\begin{aligned} SP_{x, g, k}^1 &= \{(1, 1), (x, g^3), (x, k), (1, kg^3)\}, \\ SP_{x, g, k}^2 &= \{(1, 1), (x, g^3), (1, kg), (x, kg^4)\}, \\ SP_{x, g, k}^3 &= \{(1, 1), (x, g^3), (x, kg^2), (1, kg^5)\}. \end{aligned}$$

For some  $i$  either  $RP_{x, g, h}^i$  or  $SP_{x, g, k}^i$  must also be a Sylow 2-subgroup of  $V$ . If  $V \simeq \mathbb{A}_4$ , then according to Lemma 3.6, it is of type A and hence the first components of all the elements of each of its Sylow 2-subgroups equal 1. However there are elements in  $RP_{x, g, h}^i$  and in  $SP_{x, g, k}^i$  whose first components are equal to  $x$ . So if  $V \simeq \mathbb{A}_4$ , then  $x = 1$ . Every element of  $\langle U, V \rangle$  is an alternating product of elements from  $U$  and  $V$ . Since in the semidirect product we have  $(1, t)(1, s) = (1, ts)$  for any  $t, s \in \mathbb{S}_5$ , it follows that the first component of every element from  $\langle U, V \rangle$  is 1 and hence  $\langle U, V \rangle \neq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$ .

On the other hand if  $V \simeq D_{12}$ , then according to Lemma 3.6, either  $V = R_{y, g', h'}$  or  $V = S_{y, g', k'}$  for some  $y \in \mathbb{Z}_p$ , some  $g', k' \notin \mathbb{A}_5$  and some  $h' \in \mathbb{A}_5$ . Again all the sylow 2-subgroups of  $V$  are known. The first component of each element from any sylow 2-subgroup of  $U$  is 1 or  $x$  and the first component of each element from any sylow 2-subgroup of  $V$  is 1 or  $y$ . Since  $U$  and  $V$  have at least one common Sylow 2-subgroup (namely  $U \cap V$ ), we must have  $x = y$ .

Now define  $W = (\{1\} \times \mathbb{A}_5) \cup (\{x\} \times (\mathbb{S}_5 - \mathbb{A}_5))$ . It is easy to check that  $W \leq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$ . Obviously  $U \cup V \subset W$ , and so  $\langle U, V \rangle \leq W$ . Therefore  $\langle U, V \rangle \neq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$ .  $\square$

**Proof of Theorem 3.1.** We first make a general discussion on  $G$ -semisymmetric graphs. Let  $\Gamma$  be a connected cubic  $G$ -semisymmetric graph of order  $n$ . Then  $\Gamma$  is regular and bipartite. Moreover it is  $G$ -edge-transitive and hence edge-transitive. Now if  $\Gamma$  is not vertex-transitive, then by definition it is semisymmetric cubic of order  $n$ . On the other hand if  $\Gamma$  is vertex-transitive, then it is symmetric cubic of order  $n$ , since according to [22] a cubic vertex- and edge-transitive graph is necessarily symmetric. Therefore  $\Gamma$  is either a bipartite cubic symmetric graph of order  $n$  or it is a cubic semisymmetric graph of order  $n$ .

We now set off to prove part (ii) of Theorem 3.1. For  $p = 3, 5, 7, 17, 31$  there is no connected cubic semisymmetric graph of order  $20p$  according to [5]. Also for  $p = 5, 7, 17$  no connected cubic symmetric graph of order  $20p$  exists according to [6]. As for  $p = 3, 31$ , according to [6] there exists only one connected cubic symmetric graph of order  $20p$  which is not bipartite. Therefore we conclude that for  $p = 3, 5, 7, 17, 31$  there is no connected cubic  $G$ -semisymmetric graph of order  $20p$ .

Now let  $p > 11$  be a prime such that  $p \neq 17, 31$ . Suppose on the contrary that  $\Gamma$  is a connected cubic  $G$ -semisymmetric graph of order  $20p$  for some  $G \leq \text{Aut}(\Gamma)$ . Let  $\{U, W\}$  be the bipartition for  $\Gamma$ . Then  $|U| = |W| = 10p$  and  $|G| = 2^{r+1} \cdot 3 \cdot 5 \cdot p$  for some  $0 \leq r \leq 7$ . If  $M$  is a Sylow  $p$ -subgroup of  $G$ , then according to Lemma 3.3,  $M \trianglelefteq G$ . Due to its order,  $M$  is intransitive on both  $U$  and  $W$  and so according to Theorem 2.6,  $\Gamma_M$  is a connected cubic  $G_M$ -semisymmetric graph of order 20 with the bipartition  $\{U_M, W_M\}$ , where  $G_M \simeq \frac{G}{M}$  and  $|U_M| = |W_M| = 10$ . According to the general discussion we just made,  $\Gamma_M$  is either a bipartite cubic symmetric graph or a cubic semisymmetric graph of order 20. By [5] there is no semisymmetric cubic graph of order 20 and by [6] there is only one bipartite symmetric cubic graph of order 20, namely **F20B**. Therefore  $\Gamma_M \simeq \mathbf{F20B}$ .

The automorphism group of **F20B** has 240 elements ([6]) and  $G_M$  is isomorphic to a subgroup of  $\text{Aut}(\mathbf{F20B})$  of order  $|G_M| = 2^{r+1} \cdot 3 \cdot 5$ . The equality is not possible since  $G_M$  is not transitive on  $V(\mathbf{F20B})$  whereas  $\text{Aut}(\mathbf{F20B})$  is. So  $|G_M| < 240$  and hence  $1 \leq r + 1 \leq 3$ . Also  $G_M$  is transitive on both  $U_M$  and  $W_M$  and according to Proposition 2.5, the action of  $G_M$  on each of  $U_M$  and  $W_M$  is faithful. Therefore  $G_M$  is a transitive permutation group of degree 10. Transitive permutation groups of degree 10 have been completely classified in [4]. There are 45 such groups up to isomorphism which are denoted  $T1, T2, \dots, T45$  in [4] and the only ones whose orders are of the form  $2^i \cdot 3 \cdot 5$  for  $1 \leq i \leq 3$ , are  $T7 \simeq \mathbb{A}_5$  of order 60, and  $T11, T12$  and  $T13 \simeq \mathbb{S}_5$  of order 120.

First note that  $G_M \simeq T7$  is not possible according to Lemma 3.5. Next, we argue that  $G_M$  could not be isomorphic to  $T11$  or  $T12$ .

In [4] all the transitive groups of degree 10 are defined with a set of generating permutations on ten points. If

$$a = (1, 2, 3, 4, 5), b = (6, 7, 8, 9, 10), e = (1, 5)(2, 3), f = (6, 10)(7, 8), \\ g = (1, 2), h = (6, 7) \text{ and } i = (1, 6)(2, 7)(3, 8)(4, 9)(5, 10),$$

then  $T11 = \langle ab, ef, i \rangle$  and  $T12 = \langle ab, ef, ghi \rangle$ . Using the GAP software ([23]) it is easy to verify that  $H = \langle i \rangle$  of order 2 is a normal subgroup of  $T11$ .

If  $G_M \simeq T11$ , then according to Theorem 2.6, the quotient graph of  $\Gamma_M$  with respect to  $H$  which we denote by  $(\Gamma_M)_H$ , would be  $R$ -semisymmetric of order 10, where  $R \simeq \frac{T11}{H}$ . This implies that  $R$  is transitive

1 on each partite set and by Proposition 2.5,  $R$  would be a transitive permutation group of degree 5. Again  
 2 according to [4] the only transitive permutation group of degree 5 and of order 60 is  $\mathbb{A}_5$ . So we should have  
 3  $R \simeq \mathbb{A}_5$ . Now the stabilizer of any vertex of  $(\Gamma_M)_H$  under the action of  $R$  has  $\frac{|R|}{5} = 12$  points and the  
 4 only subgroup of  $\mathbb{A}_5$  of order 12 is isomorphic to  $\mathbb{A}_4$ . So for an edge  $\{u, w\}$  of the cubic  $R$ -semisymmetric  
 5 graph  $(\Gamma_M)_H$ , we have  $(R_u, R_w) = (\mathbb{A}_4, \mathbb{A}_4)$  which is not possible according to Theorem 2.4. Therefore the  
 6 assumption that  $G_M \simeq T11$ , leads to a contradiction.

7 Now suppose  $G_M \simeq T12$ . Calculated by GAP, the stabilizer of 1 under  $T12$  is

$$8 \quad (T12)_1 = \langle (2, 4)(3, 5)(7, 9)(8, 10), (3, 5, 4)(8, 10, 9) \rangle$$

9 Again, using GAP one finds out that this group is nonabelian of order 12 which has the following group as a  
 10 normal subgroup:

$$11 \quad \langle (2, 3)(4, 5)(7, 8)(9, 10), (2, 4)(3, 5)(7, 9)(8, 10) \rangle \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$$

12 There are only 3 nonabelian groups of order 12 up to isomorphism:  $\mathbb{A}_4$ ,  $D_{12}$  and the dicyclic group of order  
 13 12. Among these, only  $\mathbb{A}_4$  has a normal subgroup of order 4. So  $(T12)_i \simeq (T12)_1 \simeq \mathbb{A}_4$  for any  $i = 1, 2, \dots, 10$ .  
 14 However this is impossible by Theorem 2.4.

15 Finally suppose  $G_M \simeq \frac{G}{M} \simeq T13$ . Since  $M \simeq \mathbb{Z}_p$ , by Lemma 3.4,  $G \simeq \mathbb{Z}_p \rtimes_{\varphi} \mathbb{S}_5$  for some homomorphism  
 16  $\varphi : \mathbb{S}_5 \rightarrow \text{Aut}(\mathbb{Z}_p)$ . From  $[G : G_u] = 10p$  we have  $|G_u| = 12$  for any vertex  $u$ . So if  $\{u, v\}$  is a fixed edge of  $\Gamma$ ,  
 17 then it follows from Theorem 2.4, that  $(G_u, G_v) \simeq (D_{12}, D_{12})$  or  $(D_{12}, \mathbb{A}_4)$ . Of course  $(G_u, G_v) \simeq (\mathbb{A}_4, D_{12})$  is  
 18 nothing new, since then we can change the roles of  $u$  and  $v$ .

19 Also  $\Gamma \simeq C(G; G_u, G_v)$  by Proposition 2.9. Now it follows from part (ii) of Proposition 2.8, that  
 20  $G = \langle G_u, G_v \rangle$  and from part (i) of the same Proposition that  $|G_u \cap G_v| = 4$ ; i.e.  $G_u \cap G_v$  is a common Sylow  
 21 2-subgroup of  $G_u$  and  $G_v$ . But the existence of  $G_u$  and  $G_v$  with all these properties contradicts Lemma 3.7.

22 Since every case for  $G_M$  is contradictory, part (ii) follows.

23 Next, we turn to part (i) of Theorem 3.1. For  $p \neq 2, 11$  there is no connected cubic semisymmetric  
 24 graph of order  $20p$  according to part (ii). Also there is no such graph of order  $20 \times 2$  according to [5] and by  
 25 the same reference, there is only one connected cubic semisymmetric graph of order  $20 \times 11$ , namely **S220**.

## 26 Acknowledgment

27 The authors would like to thank the anonymous referees for their helpful comments. The second author was  
 28 supported in part by a grant from the IMU-CDC.

## 29 References

- 30 [1] Alaeiyan M, Ghasemi M. Cubic edge-transitive graphs of order  $8p^2$ . Bulletin of the Australian Mathematical Society  
 31 2008; 77(2):315–323.
- 32 [2] Alaeiyan M, Onagh BN. On semisymmetric cubic graphs of order  $10p^3$ . Hacettepe Journal of Mathematics and  
 33 Statistics 2011; 40(4):531–535.
- 34 [3] Bugeand Y, Cao Z, Mignotte M. on simple  $K_4$ -groups. Journal of Algebra 2001; 241(2):658–668.
- 35 [4] Butler G, McKay J. The transitive groups of degree up to eleven. Communications in Algebra 1983; 11(8):863–911.

- 1 [5] Conder M, Malnič A, Marušič D, Potočnik P. A census of semisymmetric cubic graphs on up to 768 vertices. *Journal of Algebraic Combinatorics* 2006; 23(3):255–294.
- 2
- 3 [6] Conder M, Nedela R. A refined classification of symmetric cubic graphs. *Journal of Algebra* 2009; 322:722–740.
- 4 [7] Du S, Xu M. A classification of semisymmetric graphs of order  $2pq$ . *Communications in Algebra* 2000; 28(6):2685–
- 5 2715.
- 6 [8] Feng Y, Ghasemi M, Changqun W. Cubic semisymmetric graphs of order  $6p^3$ . *Discrete Mathematics* 2010;
- 7 310(17):2345–2355.
- 8 [9] Folkman J. Regular line-symmetric graphs. *Journal of Combinatorial Theory* 1967; 3(3):215–232.
- 9 [10] Goldschmidt DM. Automorphisms of trivalent graphs. *Annals of Mathematics* 1980; 111(2):377–406.
- 10 [11] Han H, Lu Z. Semisymmetric graphs of order  $6p^2$  and prime valency. *Science China Mathematics* 2012; 55(12):2579–
- 11 2592.
- 12 [12] Herzog M. On finite simple groups of order divisible by three primes only. *Journal of Algebra* 1968; 120(10):383–388.
- 13 [13] Hua X, Feng Y. Cubic semisymmetric graphs of order  $8p^3$ . *Science China Mathematics* 2011; 54(9):1937–1949.
- 14 [14] Kwak JH, Nedela R. Graphs and their coverings. *Lecture Notes Series*; vol. 17, 2007.
- 15 [15] Lu Z, Wang C, Xu M. On semisymmetric cubic graphs of order  $6p^2$ . *Science in China, Series A: Mathematics*, 2004;
- 16 47(1):1–17.
- 17 [16] Malnič A, Marušič D, Wang C. Cubic Semisymmetric Graphs of Order  $2p^3$ . Ljubljana, Slovenia: University of
- 18 Ljubljana, Preprint Series, vol. 38, 2000.
- 19 [17] Malnič A, Marušič D, Wang C. Cubic edge-transitive graphs of order  $2p^3$ . *Discrete Mathematics* 2004; 274(1-3):187–
- 20 198.
- 21 [18] Robinson DJ. *A Course in the Theory of Groups*; New York, USA: Springer-Verlag, 1982.
- 22 [19] Shi WJ. On simple  $K_4$ -groups. *Chinese Science Bulletin* 1991; 36(17):1281–1283.
- 23 [20] Suzuki M. *Group Theory*. New York, USA: Springer-Verlag, 1986.
- 24 [21] Talebi A, Mehdipoor N. Classifying cubic semisymmetric graphs of order  $18p^n$ . *Graphs and Combinatorics* 2014;
- 25 30(4):1037–1044.
- 26 [22] Tutte WT. *Connectivity in graphs*. Toronto, Canada: University of Toronto Press, 1966.
- 27 [23] The GAP Group. *Groups, Algorithms and Programming*. Version 4.8.6, 2016; <http://www.gap-system.org>.
- 28 [24] Zhang S, Shi WJ. Revisiting the number of simple  $K_4$ -groups. arXiv:1307.8079v1 [math.NT], 2013.